

# FEAST<sup>®</sup>

## MICRO MASTER<sup>™</sup>

C H E L A T E D M I C R O N U T R I E N T

- Fully chelated in a form readily available to plants
- Can be soil or foliar applied
- Use in combination with either liquid or dry fertilizers
- Compatible with most pesticides

For further information see back panel.

### KEEP FROM FREEZING

**Important:** Consult manufacturer for pertinent product information at <http://www.conklin.com/agrovan tage/docs>

## WARNING



**Harmful if swallowed or inhaled. Causes eye and skin irritation.**

**Precautions:** Avoid breathing mist, vapors and spray. Use only outdoors or in a well ventilated area. Do not eat, drink or smoke when using this product. Wear gloves, safety glasses or goggles, and long-sleeved clothing when handling. Wash hands thoroughly after handling.

9.0%  
Zinc  
EDTA Formula

**Response/First Aid:** If in eyes: Rinse cautiously with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. If eye irritation persists: Get medical advice or attention. If swallowed: Call a physician or Poison Control Center if you feel unwell. Rinse Mouth. If on skin: Wash with plenty of soap and water. If skin irritation occurs: Get medical advice or attention. Immediately take off all contaminated clothing and wash it before reuse. If inhaled: Remove person to fresh air and keep comfortable for breathing.

**Net Weight:**  
**27.5 Pounds (12.48 kg)**  
**2.5 Gallons (9.46 L)**

# FEAST<sup>®</sup>

## MICRO MASTER<sup>™</sup>

### C H E L A T E D M I C R O N U T R I E N T

**8-0-0**

**Guaranteed Analysis:**

Total Nitrogen (N) .....	8.0%
4.0% Ammonical Nitrogen	
4.0% Water Soluble Nitrogen	
Zinc (Zn) .....	9.0%
9.0% Chelated Zinc (Zn)	

Derived from Aqueous Ammonia and Zinc EDTA

Weight per gallon at 68° F.....11.0 pounds

Florida license #F048

**Notice:** This product is intended to be used to correct zinc deficiency in crops.

**Directions:** Add the proper amount of Feast<sup>®</sup> Zinc EDTA to the spray tank when it is approximately one-third full of water. Fill with water and add other chemicals, if any. (Always test for chemical compatibility prior to large-scale mixing).

**Storage and Disposal:** DO NOT reuse container. Triple rinse container prior to recycling or disposal. Keep spills from waterways and sewers.

**Limited Guarantee:** Conklin Company, Inc. warrants that this product conforms to label description. Except as expressly warranted in the preceding sentence: **DISCLAIMER OF WARRANTY: THIS LIMITED GUARANTEE IS IN LIEU OF THE IMPLIED WARRANTY OF MERCHANTABILITY, THE IMPLIED WARRANTY OF FITNESS FOR A PARTICULAR PURPOSE AND ALL OTHER WARRANTIES EXPRESS OR IMPLIED, ALL OF WHICH ARE DISCLAIMED.**

For US Health Hazard Information, call 1-888-786-0974.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapco.org/metals.htm>

**Application Guide**

**SOIL:** One to four pints per acre as needed. \* Feast<sup>®</sup> Zinc EDTA can be applied alone, with fertilizers or in combination with most pesticides. For most effective treatment, apply in combination with starter fertilizer at planting or as a sidedress shortly after.  
**NOTE:** Do not apply to organic soils such as muck or high organic soils. Instead, use a foliar application for these soil types.

**FOLIAR:** One half-pint to 2 pints per acre. When applied with ground application equipment, use no more than 10 gallons of total spray solution per acre. When applied with aerial application equipment, use no more than 5 gallons per acre of total spray solution. A quality wetting agent such as Saturall 85<sup>®</sup> or Rain-Fast<sup>®</sup> may be used to ensure even coverage. **NOTE:** if label directions are not followed for foliar applications, phytotoxicity may occur. **Foliar fertilization is intended as a supplement to a regular fertilization program and will not, by itself, provide all the nutrients normally required by agricultural crops.**

**Wisconsin Users:** Refer to supplemental sheets identifying state crops that have a medium to high zinc demand, and Wisconsin appropriate application rates.

\*Optimal rates can vary depending on soil conditions, extent of deficiency, type of crop and other factors. Conklin Company Inc. recommends the use of AgroVantage<sup>®</sup> soil and/or plant tissue analysis to accurately determine the quantities of any nutrient to be applied.

